


AD-A139 640 INTERIM REPORT FOR GRANT AFOSR-82-0146 MAY 16 1982-MAY 1/1  
15 1983(U) SYRACUSE UNIV NY SCHOOL OF COMPUTER AND  
INFORMATION SCIENCE K G MEHROTRA JUN 83  
UNCLASSIFIED AFOSR-TR-84-0177 AFOSR-82-0146 F/G 12/1 NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A139640

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

②

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 84-0177</b>	
6a. NAME OF PERFORMING ORGANIZATION Syracuse University	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) School of Computer & Information Science Syracuse, New York 13210		7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-82-0146	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332		10. SOURCE OF FUNDING NOS.	
11. TITLE (Include Security Classification) Interim Report For Grant AFOSR-82-0146, May 16, 1982-May 15, 1983		PROGRAM ELEMENT NO. 61102F	TASK NO. A5
12. PERSONAL AUTHOR(S) Kishan G. Mehrotra			
13a. TYPE OF REPORT Final	13b. TIME COVERED FROM 16 May 82 to 15 May 83	14. DATE OF REPORT (Yr., Mo., Day) Jun 1983	15. PAGE COUNT 14
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB GR.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The properties of a new rank statistic for comparing two populations in an arbitrary right censoring sampling scheme were investigated. The new statistic is an extension of the locally most powerful rank statistic for uncensored data and an extension of the well-known Wilcoxon statistic. The power of the new statistic was compared with the Prentic statistic for a sample size of 10 and the new statistic was generally of slightly higher power for the logistic alternatives investigated. The investigators are optimistic that the statistic will remain more powerful against other alternatives.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL BRIAN W. WOODRUFF, Capt, USAF		22b. TELEPHONE NUMBER (Include Area Code) (202) 767- 4927	22c. OFFICE SYMBOL

DTIC FILE COPY

DTIC  
ELECTE  
MAR 30 1984  
S  
E

84 04 03 119

FINAL REPORT.

Project Number AFOSR-82-0146

May 1982 - ~~June~~ 1983

by

Kishan G. Mehrotra

School of Computer & Information Science

Syracuse University

Syracuse, New York 13210

Phone (315) 423-2811

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



### Introduction.

In the proposal, our goal was to study rank statistics to test the hypothesis  $H_0: F^0(x)=G^0(x)$  for all  $x$ , where  $F^0$  and  $G^0$  denote the distribution functions of random variables  $X^0$  and  $Y^0$  respectively.

Let  $X_1^0, \dots, X_m^0$  and  $Y_1^0, \dots, Y_n^0$  be the two independent samples from  $F^0$  and  $G^0$  respectively, and  $N=m+n$ . Sample estimates of  $F^0(x)$  and  $G^0(x)$  are given by  $F_N^0(x)=(\# \text{ of } X_{i-}^0 \leq x)/m$  and  $G_N^0(x)=(\# \text{ of } Y_{j-}^0 \leq x)/n$  respectively, and to test  $H_0$  the locally most powerful linear rank statistic is given by

$$T_N = \int J(H_N^0(x)) dF_N^0.$$

In the above statistic,  $H_N^0 = \frac{m}{N} F_N^0(x) + \frac{n}{N} G_N^0(x)$ . The range of integration is  $(-\infty, \infty)$  here and, unless otherwise specified, in the remainder of the report.

When  $X$  and  $Y$  observations are arbitrarily right censored, a better estimator of  $F(x)$  is given by the Kaplan-Meier product limit estimator. Thus, in the case of arbitrarily right censoring situation, it was proposed to study

$$T_N^* = \int J(H_N^*(x)) dF_N^*(x),$$

where  $F_N^*$  and  $H_N^*$  denote the product limit estimator of  $F$  and  $H$  respectively. Heuristic justifications for this statistic are (AFSC)

NOTICE  
This document is approved and is  
approved for release IAW AFR 190-12.  
Distribution is unlimited.  
MATTHEW J. KERPER  
Chief, Technical Information Division

that (i) it is a natural generalization of  $T_N$ , and (ii) a generalization of the Wilcoxon statistic, as obtained by Prentice (1978) using local optimum properties, contains product limit estimators. We had proposed to study asymptotic as well as small samples properties of  $T_N^*$ .

An important special case of  $T_N$ , obtained when  $J(x)=x$ , is known as the Wilcoxon statistic. If  $J(x)=x$ ,

$$T_N = \int \left\{ \frac{m}{N} F_N^O(x) + \frac{n}{N} G_N^O(x) \right\} dF_N^O(x) = \frac{m+1}{mN} + \frac{n}{N} \int G_N^O(x) dF_N^O(x).$$

Since  $(m+1)(mN)^{-1}$  is a constant,  $T_N$  is equivalent to

$$S_N = \int G_N^O(x) dF_N^O(x).$$

The statistic  $S_N$  is attractive for yet another reason that it estimates  $\Pr[Y^O \leq X^O]$ . In the arbitrary right censoring situation, it would be appropriate to replace  $F_N^O$  and  $G_N^O$  by the Kaplan-Meier estimators  $F_N^*$  and  $G_N^*$  and therefore study

$$S_N^* = \int G_N^*(x) dF_N^*(x).$$

It turns out that Efron (1965) had obtained  $S_N^*$  as an extension of Gilbert (1962) and Gehan (1965) statistic.

In the first part of this report, Efron's statistic is compared with Prentice's statistic. In the second part, we consider some elementary properties of the proposed statistic, and in the third part, its small sample behavior is reported. It is observed that, unlike  $H_N(x)$ ,  $H_N^*(x)$  can not be expressed as

$$H_N^*(x) = \frac{m}{N} F_N^*(x) + \frac{n}{N} G_N^*(x).$$

Therefore, our statistic differs from Efron statistic. This aspect is elaborated in section 2 with its consequences.

Results of sections 2 and 3 are preliminary. Research is still continuing on some aspects of the proposed statistic. A full report will be published at the termination of such evaluations.

#### 1. Efron and Prentice Statistics.

Let  $X^0$  and  $Y^0$  be two independent random variables with distribution functions (df)  $F^0$  and  $G^0$  respectively. Let  $U$  and  $V$  be two other random variables which are independent of each other, independent of  $X^0$  and  $Y^0$ , and with corresponding df  $I_U$  and  $I_V$  respectively. In the arbitrarily right censored data, the observables are given in terms of

$$X_i = \min (X_i^0, U_i) ,$$

$$\epsilon_i = \begin{cases} 1 & \text{if } x_i^0 \leq u_i, \\ 0 & \text{otherwise,} \end{cases}$$

where  $i=1, \dots, m$ . Likewise,  $Y_j = \min(Y_j^0, V_j)$  and  $\epsilon_j = 1$  (0) if  $Y_j^0 \leq V_j$  ( $Y_j^0 > V_j$ ),  $j=1, \dots, n$  are observables from the other sample.

To compare  $F^0$  and  $G^0$ , Gehan (1965) and Gilbert (1962) independently proposed the following modification of the Wilcoxon statistic. They suggested the use of  $\sum_i \sum_j W_G(i, j)$  where the "score function"  $W_G(i, j)$  is defined as follows:

$$W_G(i, j) = \begin{cases} 1 & \text{if } x_i \geq y_j, \text{ when } y_j \text{ is an uncensored observation,} \\ 0 & \text{if } x_i < y_j, \text{ when } x_i \text{ is an uncensored observation,} \\ 1/2 & \text{if } x_i \text{ and } y_j \text{ are both censored.} \end{cases}$$

Thus, in Gehan statistic, a pair  $(x_i, y_j)$  contributes 1/2 whenever  $x_i^0$  and  $y_j^0$  "can not be compared". Efron (1965) proposed a modification to the above scoring function. He argued that the score function should be an estimate of  $P[X^0 \geq Y^0]$  and the estimate should be obtained conditional upon the available sample. This method of evaluating the score function has been shown to provide locally most powerful rank test under type II censoring by Bhattacharyya and Mehrotra (1983). Though the justification was obtained only recently, other occurrences of this conditional argument have appeared in earlier works.



We consider an example of evaluation of Efron score. If  $x_i$  is a failure,  $y_j$  is a censored observation, and  $y_j < x_i$ , then the probability of the event  $\{X^0 > Y^0\}$  is given by  $\{G^0(x_i) - G^0(y_j)\} / \{1 - G^0(y_j)\}$ . Clearly, the score associated with such a pair is

$$W_E(i, j) = \frac{G_N^*(x_i) - G_N^*(y_j)}{1 - G_N^*(y_j)},$$

where  $G_N^*$  is the product limit estimator of the distribution function  $G^0$ . As in Gehan's case,  $W_E(i, j) = 1$  if  $x_i \geq y_j$  and  $y_j$  is uncensored, 0 if  $x_i < y_j$  and  $x_i$  is uncensored. The two scoring functions differ only when the pair  $(x_i, y_j)$  can not be compared. Efron has shown that his statistic,  $T_E = \sum_i \sum_j W_E(i, j)$  can be represented as

$$T_E = \int G_N^*(x) dF_N^*(x),$$

where, as mentioned earlier,  $G_N^*$  and  $F_N^*$  are product limit estimators of  $G^0$  and  $F^0$ .

To obtain the asymptotic normality, one can apply the theory of U-statistic to the earlier form or Chernoff and Savage theorem to the integral representation of  $T_E$ . Briefly, the following result is obtained.

Theorem (Efron): Let  $m, n$  converge to infinity in such a way that  $\lim m/N = \lambda$ ,  $0 < \lambda < 1$ . Then, the distribution of

$$N^{1/2} \{T_E - \int (1-F^0) dG^0\}$$

converges to a normal distribution with mean zero and variance  $\sigma^2$ , which under the null hypothesis becomes

$$\sigma_0^2 = \frac{1}{4} \left[ \frac{1}{\lambda} \int_0^1 \frac{z^3 dz}{F\{F^{0-1}(z)\}} + \frac{1}{1-\lambda} \int_0^1 \frac{z^3 dz}{G\{F^{0-1}(z)\}} \right].$$

Here  $1-F = (1-F^0)(1-I_U)$  and  $1-G = (1-F^0)(1-I_V)$ .

From the above theorem, efficacy of Efron statistic is

$$\mathcal{E}_E = \frac{\left\{ \int (1-F^0) dG^0 \right\}^2}{\sigma_0^2}. \quad (1.1)$$

It should be remembered that the product limit estimator of the survival function  $F^0 = (1-F^0)$  is given by

$$\hat{P}(x) = \begin{cases} 1 & \text{if } x < x_{(1)}, \\ \prod_{i=1}^{k-1} \left( \frac{m-i}{m-i+1} \right)^{\delta_i} & \text{if } x_{(k)} \leq x < x_{(k+1)}, \quad k=1, \dots, m, \end{cases}$$

where  $x_{(m+1)} = \infty$ . Efron uses a slightly different estimator  $F_N^*$  of  $F^0$ . His estimator, which is self consistent i.e.

$$mF^*(s) = (\# \text{ of } x_i \geq s) + \sum_{x_i < s} (1 - \delta_i) \frac{F^*(s)}{F^*(x_i)}$$

is identical to  $\hat{P}(x)$  in all intervals  $x_{(k)} \leq x < x_{(k+1)}$  except when  $k=m$ . In this last interval  $F^*(s)=0$ ; i.e. it is assumed that  $x_{(m)}$  is uncensored, irrespective of its actual value.

Using a conditional locally most powerful criterion, Prentice (1978) obtained a rank statistic  $T_p$  for testing  $H_0$ . His statistic is given by

$$T_p = \sum c_i \{Z_{(i)} + C_i \sum_j Z_{ij}\},$$

where the outer sum is over all failures, and the inner sum is over all those observations which are censored between the  $i$ th and the  $(i+1)$ th failures;  $Z_{(i)}$  (and likewise  $Z_{ij}$ ) takes value 1 if the  $i$ th failure ( $j$ th censored) is an  $X$  observation, and value 0 otherwise. The score  $c_i$  ( $C_i$ ) corresponds to a failure (censored) observation. From the equivalence established in Mehrotra, Michalek, and Mihalko (1983), Prentice statistic can be written in a form whose asymptotic normality has been established by Shoenfeld (1982). Briefly, the efficacy of the Prentice statistic is given by

$$E_p = \frac{[\int g(t) \log\{r_2(t)/r_1(t)\} \pi(t) \{1-\pi(t)\} V(t) dt]^2}{\int g^2(t) \pi(t) \{1-\pi(t)\} V(t) dt}, \quad (1.2)$$

where

$$g(t) = \lim(c_j - C_j), \quad r_1(t) = f^0(t) / \{1 - F^0(t)\}, \quad r_2(t) = g^0(t) / \{1 - F^0(t)\},$$

$\pi(t) = (1-\lambda)\{1-G(t)\} / [\lambda\{1-F(t)\} + (1-\lambda)\{1-G(t)\}]$ , and

$$V(t) = f^0(t)\{1-I_U(t)\} + (1-\lambda)g^0(t)\{1-I_V(t)\}.$$

The ratio of (1.1) and (1.2) gives the relative efficiency of Efron statistic versus Prentice statistic.

## 2. Properties of the Proposed Statistic.

The combined ranked sample of X's and Y's can be written in terms of three vectors  $\underline{W}$ ,  $\underline{\Delta}$ , and  $\underline{Z}$ , each with dimension N. The vector  $\underline{W} = (W_1, \dots, W_N)$  represents the combined vector of ordered observations  $W_1 \leq \dots \leq W_N$ .  $\underline{\Delta}$  is a vector of indicator variables, where  $\Delta_i$  takes value 1 if  $W_i$  is a failure, and 0 otherwise.  $\underline{Z}$  is another vector of indicator variables, with  $Z_i$  taking value 1 if  $W_i$  is an X observation and 0 otherwise. Using these notations, the product limit estimator of  $H^0(x) = mN^{-1}F^0 + nN^{-1}G^0(x)$  at  $w_j$  is given by

$$H_N^*(w_j) = \prod_{i=1}^j \left( \frac{N-i}{N-i+1} \right)^{\Delta_i} = H_N^*(w_{j-1}) \left( \frac{N-j}{N-j+1} \right)^{\Delta_j}.$$

The second factor on the right hand side of the above expression, can be expressed in terms of X and Y failures as

$$\frac{N-j}{N-j+1} = \frac{(m - \sum_{k=1}^j Z_k) + (n - \sum_{k=1}^j (1-Z_k))}{(m - \sum_{k=1}^{j-1} Z_k) + (n - \sum_{k=1}^{j-1} (1-Z_k))}. \quad (2.1)$$

By convention,  $\sum_{k=1}^0 Z_k = \sum_{k=1}^0 (1-Z_k) = 0$ . On the other hand, the product limit estimators of the survival functions  $F^0$  and  $G^0$  are given by

$$F^*(w_j) = F^*(w_{j-1}) \left\{ \frac{m - \sum_{k=1}^j Z_k}{m - \sum_{k=1}^j Z_k} \right\} \Lambda_j, \quad (2.2)$$

and

$$G^*(w_j) = G^*(w_{j-1}) \left\{ \frac{n - \sum_{k=1}^j (1-Z_k)}{n - \sum_{k=1}^j (1-Z_k)} \right\} \Lambda_j. \quad (2.3)$$

From equations (2.1), (2.2), and (2.3), it is clear that, in the case of arbitrary right censoring,

$$H_N^* \neq \frac{m}{N} F_N^*(x) + \frac{n}{N} G_N^*(x).$$

As a consequence, the proposed statistic and Efron statistic differ from each other when  $J(x)=x$ . In other words, the proposed statistic is another generalization of the Wilcoxon statistic. Another interesting consequence of this observation is that, for  $J(x)=x$ , the statistic  $T_N^*$  does not estimate  $\{\text{constant} + \Pr[Y^0 \leq X^0]\}$ . This appears to be a drawback of the proposed statistic. However, weights should be assigned according to the ranks in the combined sample. This justifies the usefulness of the proposed statistic, and consequently requires further investigation.

The mass function associated with the product limit estimator  $F_N^*$  can be shown to be

$$f_N^* = \left\{ m - \frac{(1-\Lambda_1)Z_1}{F^*(w_1)} \right\}^{-1},$$

if  $w_k$  is an  $X$  observation and is uncensored. Hence, alternatively,

$$mf_N^*(w_k) = 1 + \sum_{i < k} \frac{Z_i(1-\Lambda_i)}{F^*(w_i)} f_N^*(w_k)$$

whenever  $Z_k=1$  and  $\Lambda_k=1$ . The proposed statistic  $T_N$  can be expressed as

$$T_N^* = m^{-1} \sum_{\delta_k \Lambda_k = 1} J[H_N^*(w_k)] \left\{ 1 + \frac{Z_i(1-\Lambda_i)}{F^*(w_i)} f_N^*(w_k) \right\}.$$

After changing the order of summation, the second term gives an interesting interpretation of  $T_N^*$ . Essentially, it amounts to assigning a contribution  $H_N^*(w_k)$  at each  $X$  failure. From each censored observation that falls between two failures, the contribution is the weighted sum of the mass function  $f_N^*(w_j)$ ; the weights are proportional to  $H_N^*(w_j)$  and the summation is over all future  $X$ -censored observations. Prentice statistic is similar in nature and differs in the scores associated with the censored observations.

To obtain the asymptotic normality of the statistic, we consider the following representation of  $T_N^*$ :

$$\begin{aligned} T_N^* &= \int J[H_N^*(x)] dF_N^*(x) = \int J[H^0(x)] dF^0(x) + \int J[H_N^*(x) - H^0(x)] dF^0(x) \\ &\quad + \int J[H^0(x)] d[F_N^*(x) - F^0(x)] + \int \{J[H_N^*(x)] - J[H^0(x)]\} d[F_N^*(x) - F^0(x)]. \end{aligned}$$

The first term of the above expression is a constant, and under some regularity conditions on the behavior of  $J(x)$ , the last term is asymptotically negligible. On the middle terms, one can apply

the property that  $N^{1/2}\{\hat{H}^*(s)-H^0(s)\}$ , considered as a stochastic process in  $s$ , approaches a normal process with zero mean and covariance kernel

$$H^0(s)H^0(t) \int_{-\infty}^s \frac{dH^0(z)}{\{1-H^0(z)\}\{1-H(z)\}}.$$

As a consequence, the two middle terms are asymptotically normally distributed with zero mean and appropriately obtained variance.

These and other related details are still under further investigations.

### 3. Small Sample Behavior of $T_N^*$ .

At the present time, we have investigated the behavior of  $T_N^*$  for  $J(x)=x$  which provides a generalized Wilcoxon statistic. We have compared this statistic with Prentice statistic which, in this section, will be denoted by  $T_p$ . This comparison is made when the  $X$  and  $Y$  observations are generated from logistic populations. Prentice statistic is obtained with the appropriate "logistic" weights. Censoring varies over 0%, 10%, 30%, and 50%. The zero percent censoring is used to check the accuracy of the simulation results. Clearly, in this case,  $T_N^*$  and  $T_p$  are essentially the same, and both equivalent to the Wilcoxon statistic. In the alternative situation, the censored  $Y$  observations are generated by varying the location parameter,  $\beta$ , of the logistic

distribution from 0.1 to 0.9. In every case, the censoring distribution is a uniform whose range is chosen so that the desired censoring probability is attained. The sample sizes of the populations X and Y are both kept equal to 10. The power is obtained from 1000 repetitions. Table 1 shows 1000xpower of the statistics  $T_N^*$  and  $T_P$ .

TABLE 1: Simulated Powers of the Prentice and Proposed Statistics.

Prob. of Censor.	0.0		0.1		0.3		0.5	
$\beta$	$T_P$	$T_N^*$	$T_P$	$T_N^*$	$T_P$	$T_N^*$	$T_P$	$T_N^*$
0.0	50	50	50	50	50	50	50	50
0.1	56	56	60	65	47	57	55	76
0.2	59	59	81	82	55	69	65	80
0.3	84	84	96	106	65	78	71	79
0.4	109	109	115	121	99	119	107	113
0.5	148	148	151	165	136	134	133	151
0.6	167	167	179	195	147	171	147	156
0.7	196	196	185	190	175	173	160	146
0.8	225	225	236	230	189	188	192	179
0.9	292	292	285	296	225	227	180	190

Our simulation results show that the power of the proposed statistic  $T_N^*$  is generally larger than the power of  $T_P$ , though the difference is relatively small. This leads us to believe that  $T_N^*$  will continue to perform at least as well as Prentice statistic, even if the X's and Y's are generated from other distributions. Of course, the J function in  $T_N^*$  and the scores in the Prentice statistic must be chosen appropriately.



A simulation study, covering a wider range of distributions and sample sizes, is in progress. A technical report, based on this large study, is expected in the near future.

#### References.

Bhattacharyya, Gouri K. and Mehrotra, Kishan G. (1983), "A unified treatment of locally most powerful rank tests under type II censoring," Journal of the American Statistical Association, 78, 375-381.

Chernoff, H. and Savage, R. (1958), "Asymptotic normality and efficiency of certain nonparametric test statistics," Annals of Mathematical Statistics, 29, 972-994.

Gehan, E.A. (1965), "A generalized Wilcoxon test for comparing arbitrarily singly-censored samples," Biometrika, 52, 203-223.

Efron, Bradley (1967), "On two sample problem with censored data," Proceeding of the Fifth Berkeley Symposium in Mathematical Statistics, IV, Prentice Hall, New York.

Mehrotra, Kishan G., Michalek, Joel E., and Michalko, Daniel (1981), "A comparative study of some linear rank tests for the two sample problem subject to right censoring," Technical Report, School of Computer and Information Science, Syracuse University, Syracuse, N.Y. 13210.

Prentice, R.L. (1978), "Linear rank tests with right censored data," Biometrika, 65, 267-279.

Schoenfeld, D. (1981), "The asymptotic properties of nonparametric tests for comparing survival distributions," Biometrika, 68, 316-319.

END

FILMED

5-84

DTIC